

<222> (1)..(1)

87534-3000.txt SEQUENCE LISTING

	7 3 13
<110>	Hornik, Vered
<120>	CONFORMATIONALLY CONSTRAINED BACKBONE CYCLIZED SOMATOSTATIN ANALOGS
<130>	87534-3000
<140>	09/734,583
<141>	2000-12-13
<160>	10
<170>	PatentIn version 3.1
<210>	1
<211>	14
<212>	PRT
<213>	mammalian
<400>	1
Ala Gly 1	y Cys Lys Asn Phe Phe Trp Lys Thr Phe Thr Ser Cys 5 10
<210>	2
<211>	6
<212>	PRT
<213>	Artificial sequence
<220>	
<221>	DISULFIDE BRIDGE

<223> Cys residues at amino acid positions and 6 form a disulfide bridge

```
<220>
<221> MOD_RES
<222> (3)..(3)
<223>
     The Trp residue is the D isomer
<220>
<223>
      Synthetic peptide
<400> 2
Cys Phe Trp Lys Thr Cys 5
<210> 3
<211> 6
<212> PRT
<213> Artificial peptide
<220>
<221> MOD_RES
<222> (1)..(1)
<223> N-Methyl
<220>
<221> MOD_RES
<222> (1)..(6)
<223> cyclo
<220>
<221> MOD_RES
<222> (3)..(3)
<223> The Trp residue is the D isomer
<220>
<223> Synthetic peptide
<400> 3
Ala Tyr Trp Lys Val Phe
1 5
```

<210> 4

```
<211> 8
<212> PRT
<213> Artificial Sequence
<220>
<221> MOD_RES
<222> (1)..(1)
<223> The Phe residue is a D isomer
<220>
<221> MOD_RES
<222> (8)..(8)
<223> The Thr residue ends with CH2OH
<220>
<221> DISULFIDE BRIDGE
<222> (2)..(2)
<223> A disulfide bride is formed between Cys residues 2 and 7
<220>
<221> MOD_RES
<222> (4)..(4)
<223> The Trp residue is a D isomer
<220>
<223> Synthetic peptide
<400> 4
Phe Cys Phe Trp Lys Thr Cys Thr 1
<210> 5
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
```

<221> DISULFIDE

- <222> (2)..(2)
- A Disulfide Bridge is formed between the Cys residues at position <223> 2 and 6
- <220>
- <221> MOD_RES
- <222> (1)..(1)
- <223> The Phe residue is a D isomer
- <220>
- <221> MOD_RES
- <222> (4)..(4)
- <223> The Trp residue is a D isomer
- <220>
- <221> MOD_RES
- <222> (7)..(7)
- <223> The Thr resiude ends with N2H
- <220> <223> Synthetic peptide
- <400> 5
- Phe Cys Phe Trp Lys Cys Thr
- <210> 6
- <211> 8
- <212> PRT
- <213> Artificial sequence
- <220>
- <221> MISC_FEATURE
- <222> (1)..(1)
- is a gamma amino butyric acid, diamino butyric acid, Gly, beta-Al a, 5-amino pentanoic acid or amino hexanoic acid; Residue 1 is bridged to Residue 8; Residue 1 also begins with a hydrogen, or a mono- or di- saccharide attached <223>
- <220>
- <221> MISC_FEATURE
- <222> (2)..(2)

87534-3000.txt

```
<223>
       is (D) or (L) Phe or Tyr
<220>
<221>
        MISC_FEATURE
<222>
        (3)..(3)
        is (D) or (L)-Trp, or (L)-Phe, (D)- or (L)-1Nal or (D) or (L)-2Nal, or Tyr \,
<223>
<220>
<221>
        MISC_FEATURE
<222>
        (4)..(4)
<223>
        is (D) or (L)-Trp
<220>
<221>
        MISC_FEATURE
<222>
        (5)..(5)
<223>
        is (D) or (L)-Lys
<220>
<221>
        MISC_FEATURE
<222>
        (6)..(6)
        is Thr, Gly, Abu, Ser, Cys, Val, (D) or (L)-Ala, or (D)- or (L)-Ala, or Tyr \,
<223>
<220>
<221>
        MISC_FEATURE
<222>
        (7)..(7)
        is (D) or (L)-Phe, or (D)- or (L)-Ala, Nle, or Cys
<223>
<220>
<221>
        MISC_FEATURE
<222> (8)..(8)
        is Gly, Val, Leu, (D) or (L)-Phe, or 1Nal or 2Nal; with a terminal carboxy acid, amide or alcohol group
<223>
<220>
<223> Synthetic peptide
<400> 6
xaa xaa xaa xaa xaa xaa xaa 1
```

```
<210> 7
<211>
       7
<212>
       PRT
<213> Artificial Sequence
<220>
<221>
       MISC_FEATURE
<222>
        (1)..(1)
        is (D)- or (L)-Phe, or (D)- or (L)-Ala; wherein Residue 1 is bridged to Residue 6 a bridging group composed of 1 to 5 methyl spacers connected to an amide, thioether, thioester, or disulfide, followed by 1 to 5 methyl spacers \frac{1}{2}
<223>
<220>
<221>
        MISC_FEATURE
<222>
         (2)..(2)
<223>
        is Tyr or (D)- or (L)-Phe
<220>
<221> MISC_FEATURE
<222>
        (3)..(3)
        is (D)- or (L)-Trp, (D)- or (L)-1Nal, or (D)- or (L)-2Nal
<223>
<220>
<221> MISC_FEATURE
<222>
        (5)..(5)
<223>
        is Thr, Val, Ser, or Cys
<220>
<221>
        MISC_FEATURE
<222>
         (6)..(6)
<223>
        is Gly or (D)- or (L)-Phe
<220>
<221> MISC_FEATURE
<222>
        (7)..(7)
        is Thr, GABA, (D)- or (L)-1Nal, (D)- or (L)-2Nal, or (D)- or
<223>
         (L)-Phe
```

<220>

<223> Synthetic peptide

```
<400> 7
Xaa Xaa Xaa Lys Xaa Xaa Xaa
1 5
<210> 8
<211> 9
<212> PRT
<213> Artificial Sequence
<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> is absent or is any amino acid
<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> is absent or is any amino acid
<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> is absent or is any amino acid
<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> is absent or is any amino acid
<220>
<221> MISC_FEATURE
<222>
      (5)..(5)
      is 1Nal, 2Nal, Beta-Asp (Ind), Gly, Tyr, (D)- or (L)-Ala, or (D)- or (L)-Phe \,
<223>
<220>
<221> MISC_FEATURE
```

<222> (6)..(7)

```
87534-3000.txt
       may be absent, or are independently Gly, Tyr, 1Nal, 2Nal, Beta-Asp (Ind), Gly, Tyr, (D)- or (L)-Ala, or (D)- or (L)-Phe
<220>
<221>
       MISC_FEATURE
<222>
       (8)..(8)
<223>
       (D) - or (L) -Trp
<220>
<221>
       MISC_FEATURE
<222>
       (9)..(9)
       (D)- or (L)-Lys
<223>
<220>
<221>
       MISC_FEATURE
<222>
       (10)..(10)
       is absent or is Gly, Abu, Cys, Thr, Val, (D)- or (L)-Ala, or (D)- or (L)-Phe \,
<223>
<220>
<221>
       MISC_FEATURE
<222>
       (11)..(11)
       is Cys, (D)- or (L)-Ala, or (D)- or (L)-Phe
<223>
<220>
<221>
       MISC_FEATURE
<222>
       (12)..(12)
       is absent or is Val, Thr, 1Nal or 2Nal
<223>
<220>
       Synthetic peptide
<223>
<400> 8
<210>
      9
<211> 7
<212>
       PRT
<213> Artificial Sequence
```

```
87534-3000.txt
<220>
        MISC_FEATURE
<221>
<222>
         (1)..(1)
         (D)- or (L)-Phe, Tyr or (D)- or (L)-Ala; Residue 1 is connected to Residue 7 by a bridge comprised of 1 to 5 methylene spacers connected to an amide, thioether, thioester, or disulfide, followed by 1 to 5 methylene spacers
<223>
<220>
<221>
        MISC_FEATURE
<222>
         (2)..(2)
<223>
         (D)- or (L)-Phe, Tyr or (D)- or (L)-Ala;
<220>
<221>
        MISC_FEATURE
<222>
         (3)..(3)
        is absent or is (D)- or (L)-Phe, Tyr or (D)- or (L)-Ala
<223>
<220>
<221>
        MISC_FEATURE
<222>
         (4)..(4)
<223>
        is (D)- or (L)Tyr
<220>
<221> MISC_FEATURE
<222>
         (5)..(5)
<223>
        is (D)- or (L)-Lys
<220>
<221>
        MISC_FEATURE
<222>
         (6)..(6)
         is absent or is Thr, Val, Cys or (D)- or (L)-Ala
<223>
<220>
<221> MISC_FEATURE
<222>
         (7)..(7)
```

is a (D)- or (L)-Phe, Cys, or (D)- or (L)-Ala

<223> <220> <223>

<400>

Synthetic peptide

```
Xaa Xaa Xaa Xaa Xaa Xaa 1
```

- <210> 10
- <211> 7
- <212> PRT
- <213> Artificial Sequence
- <220>
- <221> MISC_FEATURE
- <222> (1)..(1)
- <223> is absent or is (D)- or (L)-Phe or Ala; the bride is connected to Residue 1 or 2 and Residue 6 or 7, wherein the birdge is comprised of 1 to 5 methylene spacers connected to an amide, thioether, thioester, or disulfide, followed by 1 to 5 methylene spacers
- <220>
- <221> MISC_FEATURE
- <222> (2)..(2)
- <223> is (D)- or (L)-Phe or Ala, Tyr
- <220>
- <221> MISC_FEATURE
- <222> (3)..(3)
- <223> is (D)- or (L)-Trp
- <220>
- <221> MISC_FEATURE
- <222> (4)..(4)
- <223> is (D)- or (L)-Lys
- <220>
- <221> MISC_FEATURE
- <222> (5)..(5)
- <223> is Thr, Ala, Val, or Cys
- <220>
- <221> MISC_FEATURE

- <222> (6)..(6)
- <223> is absent or is (D)- or (L)-Phe, Ala, or Cys
- <220>
- <221> MISC_FEATURE
- <222> (7)..(7)
- <223> is absent or is Thr or Thr reduced to an alcohol
- <220> <223> Synthetic peptide
- <400> 10
- Xaa Xaa Xaa Xaa Xaa Xaa 1